

DRUG FACTS

Active ingredient(s)

Purpose

Use(s)

- temporarily relieves minor aches and pains due to:
- minor pain of arthritis
- muscular aches
- backache
- premenstrual and menstrual cramps
- the common cold
- headache
- toothache
- temporarily reduces fever

Warnings

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have difficulty swallowing large tablets or capsules. People over 65 may have difficulty swallowing these tablets.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product

Stop use and ask a doctor if

- the tablet got stuck in your throat
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding

Keep out of reach of children

Keep out of the reach of children.

Directions

- do not take more than directed (see overdose warning)

adults	<ul style="list-style-type: none">take 2 gelcaps every 8 hours. Swallow only one gelcap at a time.take a sip of water before swallowing each gelcap and wash each gelcap down with water (up to a full 8 oz glass)swallow whole – do not crush, chew, split or dissolvedo not take more than 6 gelcaps in 24 hoursdo not use for more than 10 days unless directed by a doctor
under 18 years of age	<ul style="list-style-type: none">ask a doctor

Other information

- store at 20–25°C (68–77°F). Avoid high humidity.
- do not use if carton is opened or neck wrap or foil inner seal imprinted with "Safety Seal®" is broken
- see end panel for lot number and expiration date

Drug Facts

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 6 gelcaps in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Inactive ingredients

benzyl alcohol, butylparaben, castor oil, corn starch, edetate calcium disodium, FD&C blue #1 aluminum lake, FD&C blue #2 aluminum lake, gelatin, hydroxyethyl cellulose, hypromellose, magnesium stearate, methylparaben, microcrystalline cellulose, povidone, powdered cellulose, pregelatinized starch, propylene glycol, propylparaben, shellac, sodium lauryl sulfate, sodium propionate, sodium starch glycolate, titanium dioxide